

Technical Data Sheet

Biotin Mouse Anti-Human CD123

Product Information

Material Number:	554528
Alternate Name:	IL-3 Receptor α chain
Size:	0.5 mg
Concentration:	0.5 mg/ml
Clone:	7G3
Immunogen:	Human IL-3Ra-transfected cells
Isotype:	Mouse IgG2a, κ
Reactivity:	QC Testing: Human
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

Description

Reacts with human CD123, the 70 kD IL-3 receptor α chain (IL-3R α), which associates with the 120-140 kD β subunit. The β chain is shared with the receptors for interleukins IL-5 and GM-CSF. IL-3R α is expressed on hematopoietic progenitors and plays an important role in hematopoietic progenitor cell growth and differentiation. This antibody has been reported to block the binding of 125I-IL-3 to high and low affinity IL-3 receptors. In functional experiments, this antibody was found to inhibit acute myeloid leukemia cell proliferation, basophil histamine release, endothelial cell-mediated IL-8 secretion, and neutrophil transmigration. This antibody has been reported to be useful for immunoprecipitation, western blot and immunofluorescent staining for flow cytometry. At the Fifth HLDA Workshop, the human IL-3 receptor was designated CD123.

This antibody is routinely tested by flow cytometric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with biotin under optimum conditions, and unreacted biotin was removed.

Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application

Flow cytometry	Routinely Tested
----------------	------------------

Recommended Assay Procedure:

Immunofluorescent Staining and Flow Cytometric Analysis: The 7G3 antibody is useful for immunofluorescent staining and flow cytometric analysis to identify and enumerate CD123 producing cells within mixed cell populations and can be used with Streptavidin PE (Cat. No. 554061). The purified format of this antibody can be used in conjunction with biotinylated goat anti-mouse Ig with no cross reactivity to human Ig and streptavidin-phycoerythrin in a three layer staining procedure to amplify immunofluorescent signals for the flow cytometric analysis of cytokine receptor-expressing cells.

Neutralization: The NA/LE™ format of the 7G3 antibody (Cat. No. 554526) can be used to neutralize IL-3 bioactivities.

Suggested Companion Products

Catalog Number	Name	Size	Clone
554061	PE Streptavidin	0.5 mg	(none)

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2006 BD



BD

BD Biosciences

References

- Schlossman SF, Boumsell L, Gilks W, et al, ed. *Leukocyte Typing V: White Cell Differentiation Antigens*. New York: Oxford University Press; 1995.(Clone-specific)
- Korpelainen EI, Gamble JR, Smith WB, et al. The receptor for interleukin 3 is selectively induced in human endothelial cells by tumor necrosis factor alpha and potentiates interleukin 8 secretion and neutrophil transmigration. *Proc Natl Acad Sci U S A*. 1993; 90(23):11137-11141.(Clone-specific)
- Sun Q, Woodcock JM, Rapoport A, et al. Monoclonal antibody 7G3 recognizes the N-terminal domain of the human interleukin-3 (IL-3) receptor alpha-chain and functions as a specific IL-3 receptor antagonist. *Blood*. 1996; 87(1):83-92.(Immunogen: Blocking, Immunoprecipitation, Neutralization, Western blot)
- Zola H. Detection of cytokine receptors by flow cytometry. In: Coligan JE, Kruisbeek AM, Margulies DH, Shevach EM, Strober W, ed. *Current Protocols in Immunology*. New York: Green Publishing Associates and Wiley-Interscience; 1995:6.21.1-6.21.18.(Clone-specific: Flow cytometry)