

Product Data Sheet

Biotin anti-human CD158e1 (KIR3DL1, NKB1)

Catalog # / Size: 2163520 / 100 µg

Clone: DX9

Isotype: Mouse IgG1, κ

Immunogen: Human NK cell clone VL186-1.6


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with biotin under optimal conditions. The solution is free of unconjugated biotin.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted between 2°C and 8°C. Do not freeze.



Human peripheral blood lymphocytes stained with DX9 PE and CD16 FITC

Applications:

Applications: FC - *Quality tested*

Application Notes: The DX9 antibody reacts with the KIR (killer cell inhibitory receptor) designated NKB1 or KIR3DL1. Additional reported applications (for the relevant formats) include: immunoprecipitation¹ and restoring the NK cell cytotoxicity^{4,8}.

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 0.5 µg per 10⁶ cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application References:

1. Litwin V, *et al.* 1994. *J. Exp. Med.* 180:537. (IP)
2. Gumperz J, *et al.* 1996. *J. Exp. Med.* 183:1817.
3. Gardiner CM, *et al.* 2001. *J. Immunol.* 166:2992.
4. Bakker ABH, *et al.* 1998. *J. Immunol.* 160:5239.
5. Goodier M, *et al.* 2000. *J. Immunol.* 165:139.
6. Kirwan SE and Burshtyn DN. 2005. *J. Immunol.* 175:5006. (FC)
7. Yawata M, *et al.* 2002. *Immunogenetics* 54:543.
8. Valiante NM, *et al.* 1997. *Immunity* 7:739.
9. Pascal V, *et al.* 2007. *J. Immunol.* 179:1625. (FC) PubMed
10. Lichterfeld M, *et al.* 2008. *J. Exp. Med.* 204:2813. (FC) PubMed

Description: CD158e1, also known as NKB1, is a 70 kD member of the immunoglobulin superfamily that is expressed on a subset of natural killer cells and T cells at varying levels among individuals. NKB1 is a type I membrane protein containing two immunoglobulin C2-type domains. The interaction of NKB1 with specific HLA-B antigens on a target cell (the HLA-Bw4 allele, for example) inhibits cytotoxicity and prevents target cell lysis and death. The interactions between KIR and MHC class I are thought to be important in NK and T cell regulation following antigen stimulation. The absence of ligands for KIRs may lower the threshold for activation through activating receptors and increase inflammation and susceptibility to autoimmune disease.

Antigen References:

1. Colonna M, *et al.* 1995. *Science* 268:405.
2. D'Andrea A, *et al.* 1995. *J. Immunol.* 155:2306.
3. Uhrburg M, *et al.* 1997. *Immunity* 7:753.
4. Gumperz JE, *et al.* 1996. *J. Exp. Med.* 183:1817.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.

5. Wagtmann N, *et al.* 1995. *Immunity* 3:801.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.

Sony Biotechnology Inc. 2100 South Oak Street, Champaign, IL 61820-0903 www.sonybiotechnology.com
Toll-Free Phone: 1-(800) 275-5963 Phone: (217) 328-9396 Fax: (217) 328-9692