Product Data Sheet

PerCP/Cy5.5 anti-human CD205 (DEC-205)

Catalog # / Size: 2311045 / 25 tests

2311050 / 100 tests

Clone: HD30

Isotype: Mouse IgG1, κ

Reactivity: Human

Preparation: The antibody was purified by affinity

chromatography, and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5


and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing

0.09% sodium azide and 0.2% (w/v) BSA (origin

Storage: The antibody solution should be stored undiluted

between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.



Human monocyte-derived dendritic cells stained with HD30 PerCP/Cy5.5

Applications:

Applications: FC - Quality tested

Application Notes: Additional reported applications (for the relevant formats) include:

immunohistochemical staining of fixed, frozen sections⁴ and Western blotting

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent

staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells or 5 µl per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal

performance for each application.

* PerCP/Cy5.5 has a maximum absorption of 482 nm and 564 nm and a

maximum emission of 690 nm.

Application References: 1. Guo M, et al. 2000. Human Immunology 61:729.

2. Cho H, et al. 2007. Physiol Genomics doi:10.1152/physiolgenomics.00051.2006

3. Gurer C, et al. 2008. Blood 112:1231. PubMed

4. Park CG, et al. 2012. J. Immunol. Methods 377:15. (IHC, WB)

Description: CD205 is a 210 kD C-type lectin transmembrane protein, known as DEC-205.

It belongs to macrophage mannose receptor family and is found at high levels on dendritic cells and thymic epithelial cells. Unlike murine CD205, human CD205 is also expressed at low levels on T- and B-cells, NK cells and monocytes. CD205, serves as an endocytic receptor, functions in antigen

uptake/processing and clearance of apoptotic cells.

Antigen References: 1. Guo M, et al. 2000. Human Immunology 61:729.

2. Kato M, et al. 2006. Int. Immunol.

3. Kato M, *et al.* 2003. *J. Biol. Chem.* 278(36):34035. 4. Small M and G. Kraal. 2003. *Int. Immunol.* 15(2):197.

5. Mahnke K, et al. 2000. J. Cell Biol. 151(3):673.

This product or portions thereof is manufactured under license from GE Healthcare, under U.S. Patent Numbers 5,268,486: 5,569,587: 5,627,027 and patents or pending applications that are continuations. continuations-in-part, re-examinations, divisionals, reissues or foreign equivalents thereof. THIS MATERIAL IS SUBJECT TO PROPRIETARY RIGHTS OF GE HEALTHCARE BIO-SCIENCES CORP. AND CARNEGIE MELLON UNIVERSITY AND MADE AND SOLD UNDER LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. THIS PRODUCT IS LICENSED FOR SALE ONLY FOR RESEARCH. IT IS NOT LICENSED FOR ANY OTHER USE. THERE IS NO IMPLIED LICENSE HEREUNDER FOR ANY COMMERCIAL USE.

COMMERCIAL USE shall include: 1. sale, lease, license or other transfer of the material or any material derived or produced from it; 2. sale, lease, license or other grant of rights to use this Material or any material derived or produced from it; 3. use of this material to perform services for a fee for third parties. IF YOU REQUIRE A COMMERCIAL LICENSE TO USE THIS MATERIAL AND DO NOT HAVE ONE RETURN THIS MATERIAL, UNOPENED TO Sony Biotechnology Inc., 2100 South Oak Street, Champaign, IL 6182:
USA AND ANY MONEY PAID FOR THE MATERIAL WILL BE REFUNDED.
For recovery was only. Not for disconnections. Not for recola. Comp. Distochards and the will not be held recognible.