

Biotin anti-human CD11c

Catalog # / Size: 2108055 / 25 µg
2108060 / 100 µg

Clone: 3.9

Isotype: Mouse IgG1, κ

Workshop Number: III NL707


Reactivity: Human, **Cross-Reactivity:** Chimpanzee, Baboon, African Green, Cynomolgus, Rhesus, Squirrel Monkey

Preparation: The antibody was purified by affinity chromatography, and conjugated with biotin under optimal conditions. The solution is free of unconjugated biotin.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted between 2°C and 8°C. **Do not freeze.**



Human peripheral blood granulocytes stained with biotinylated 3.9, followed by Sav-PE

Applications:

Applications: FC - Quality tested

Application Notes: Clone 3.9 preferentially binds the activated form of CD11c, is specific for I domain of CD11c, and is able to partially block the binding of CD11c and ICAM-4. 3.9 binding is divalent cation dependent.¹² While analyzing blood, it is best to use heparin as the anti-coagulant and not EDTA. Since the ability of clone 3.9 to bind to its target is divalent cation dependent, the usage of EDTA as an anti-coagulant may be detrimental to staining due to its chelating properties.

Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections⁴, and functional assays^{5,6}.

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤2.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application References:

- Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
- Knapp W, *et al.* 1989. Leucocyte Typing IV Oxford University Press. New York.
- McMichael A, *et al.* Eds. 1987. Leucocyte Typing III Oxford University Press. New York.
- Vainer B, *et al.* 2000. *Am. J. Surg. Pathol.* 24:1115. (IHC)
- Ottonello L, *et al.* 1999. *Blood* 93:3505.
- Metelitsa LS, *et al.* 2002. *Blood* 99:4166.
- Sadhu C, *et al.* 2007. *J. Leukoc. Biol.* doi:10.1189/jlb.1106680. PubMed
- Ihanus E, *et al.* 2007. *Blood* 109:802-810.
- Gurer C, *et al.* 2008. *Blood* 112:1231. PubMed
- Asai A, *et al.* 2009. *J. Lipid Res.* 50:95. PubMed
- Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
- Sadhu C, *et al.* 2008. *J. Immunoass. Immunoch.* 29:42. (FC)

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.

Description: CD11c is a 145-150 kD type I transmembrane glycoprotein also known as integrin α_X and CR4. CD11c non-covalently associates with integrin β_2 (CD18) and is expressed on monocytes/macrophages, dendritic cells, granulocytes, NK cells, and subsets of T and B cells. CD11c has been reported to play a role in adhesion and CTL killing through its interactions with fibrinogen, CD54, and iC3b.

Antigen References:

1. Petty H. 1996. *Immunol. Today* 17:209.
2. Springer T. 1994. *Cell* 76:301.
3. Ihanus E, *et al.* 2007. *Blood* 109:802-810.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.

Sony Biotechnology Inc. 2100 South Oak Street, Champaign, IL 61820-0903 www.sonybiotechnology.com
Toll-Free Phone: 1-(800) 275-5963 Phone: (217) 328-9396 Fax: (217) 328-9692