


Brilliant Violet 711™ anti-mouse/human CD45R/B220

Catalog # / Size: 1116275 / 50 µg
Clone: RA3-6B2
Isotype: Rat IgG2a, κ
Immunogen: Abelson murine leukemia virus-induced pre-B tumor cells
Reactivity: Human
Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 711™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 711™ and unconjugated antibody.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).
Concentration: 0.2



C57BL/6 mouse splenocytes were stained with CD45R/B220 (clone RA3-6B2) Brilliant Violet 711™ (filled histogram) or rat IgG2a, κ Brilliant Violet 711™ isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤0.25 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 711™ excites at 405 nm and emits at 711 nm. The bandpass filter 710/50 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. **Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel.** Refer to your instrument manual or manufacturer for support. Brilliant Violet 711™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: Clone RA3-6B2 has been described to react with an epitope on the extracellular domain of the transmembrane CD45 glycoprotein which is dependent upon the expression of exon A and specific carbohydrate residues. Additional reported applications (for the relevant formats) include: immunoprecipitation¹, *in vitro* and *in vivo* modulation of B cell responses²⁻⁴, and immunohistochemistry of acetone-fixed frozen sections and formalin-fixed paraffin-embedded sections^{5,6}. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 103216).

Application 1. Coffman RL. 1982. *Immunol. Rev.* 69:5. (IP)

- References:**
2. George A, *et al.* 1994. *J. Immunol.* 152:1014. (Activ)
 3. Asensi V, *et al.* 1989. *Immunology* 68:204. (Activ)
 4. Domiati-Saad R, *et al.* 1993. *J. Immunol.* 151:5936. (Activ)
 5. Hata H, *et al.* 2004. *J. Clin. Invest.* 114:582. (IHC)
 6. Monteith CE, *et al.* 1996. *Can. J. Vet. Res.* 60:193. (IHC)
 7. Shih FF, *et al.* 2006. *J. Immunol.* 176:3438. (FC)
 8. Chang C L-T, *et al.* 2007. *J. Immunol.* 178:6984.
 9. Fazilleau N, *et al.* 2007. *Nature Immunol.* 8:753.
 10. Lang GL, *et al.* 2008. *Blood* 111:2158. [PubMed](#)
 11. Charles N, *et al.* 2010. *Nat. Med.* 16:701. (FC) [PubMed](#)
 12. del Rio ML, *et al.* 2011. *Transpl. Int.* 24:501. (FC) [PubMed](#)
 13. Murakami R, *et al.* 2013. *PLoS One.* 8:73270. [PubMed](#)
-

Description: CD45R, also known as B220, is an isoform of CD45. It is a member of the protein tyrosine phosphatase (PTP) family with a molecular weight of approximately 180-240 kD. CD45R is expressed on B cells (at all developmental stages from pro-B cells through mature B cells), activated B cells, and subsets of T and NK cells. CD45R (B220) is also expressed on a subset of abnormal T cells involved in the pathogenesis of systemic autoimmunity in MRL-*Fas*^{lpr} and MRL-*Fas*^{gld} mice. It plays a critical role in TCR and BCR signaling. The primary ligands for CD45 are galectin-1, CD2, CD3, and CD4. CD45R is commonly used as a pan-B cell marker; however, CD19 may be more appropriate for B cell specificity.

- Antigen**
- References:**
1. Barclay A, *et al.* 1997. The Leukocyte Antigen FactsBook Academic Press.
 2. Trowbridge IS, *et al.* 1993. *Annu. Rev. Immunol.* 12:85.
 3. Kishihara K, *et al.* 1993. *Cell* 74:143.
 4. Pulido R, <