#### Brilliant Violet 421™ anti-human IL-2

Catalog # / Size: 3101635 / 25 tests

3101640 / 100 tests

Clone: MQ1-17H12 Isotype: Rat IgG2a, κ

**Immunogen:** E. coli - expressed recombinant human

IL-2

Reactivity: Human

**Preparation:** The antibody was purified by affinity

chromatography and conjugated with Brilliant Violet 421  $^{\text{\tiny M}}$  under optimal conditions. The solution is free of unconjugated Brilliant Violet 421  $^{\text{\tiny M}}$  and


unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide and BSA

(origin USA).

Concentration: Lot-specific



Human peripheral blood lymphocytes were stimulated with PMA + Ionomycin for 6 hours (in the presence of monensin), surface stained with CD3 FITC, fixed, permeabilized, and then stained with IL-2 (clone MQ1-17H12) Brilliant Violet 421™ (top) or rat I

#### **Applications:**

**Applications:** Flow Cytometry

Recommended

**Usage:** 


Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be

each application.

Brilliant Violet 421<sup>™</sup> excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421<sup>™</sup> is a trademark of Sirigen Group Ltd.

titrated for optimal performance for

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by



U.S. Patent(s), pending patent applications and foreign equivalents.

### Application Notes:

ELISA or ELISPOT Capture<sup>2,3</sup>: The purified MQ1-17H12 antibody is useful as the capture antibody in a sandwich ELISA or ELISPOT assay, when used in conjunction with the biotinylated Poly5176 antibody (Cat. No. 517605) as the detecting antibody. The LEAF™ purified antibody is suggested for ELISPOT capture. For ELISPOT capture applications, a concentration range of 4-8 microg/ml is recommended.

Additional reported applications (for the relevant formats) include:

immunoprecipitation2, immunohistochemical staining of paraformaldehyde-fixed, saponintreated frozen tissue sections<sup>1,4-6,8</sup>, neutralization<sup>13</sup>, and immunocytochemistry.

**Note:** For testing human IL-2 in serum or plasma, BioLegend's LEGEND MAX™ Kits (Cat. No. 431807 & 431808) are specially developed and recommended.

## Application References:

- 1. Andersson J, et al. 1994. Immunology 83:16. (IHC)
- 2. Abrams J, et al. 1992. Immunol. Rev. 127:5. (IP)
- 3. Abrams JS. 1995. Curr. Prot. Immunol. Unit 6.20.
- 4. Fernandez V, et al. 1994. Eur. J. Immunol. 24:1808. (IHC)
- 5. Skansen-Saphir U, et al. 1994. Eur. J. Immunol. 24:916. (IHC)
- 6. Andersson U, et al. Detection and Quantification of Gene Expression. New York:Springer-Verlag. (IHC)
- 7. Prussin C, et al. 1995. J. Immunol. Methods. 188:117.
- 8. Raqib R, et al. 2002. Infect. Immun. 70:3199. (IHC)
- 9. Dzhagalov I, et al. 2007. J. Immunol. 178:2113. PubMed
- 10. Colleton BA, et al. 2009. J Virol. 83:6288. PubMed
- 11. Yoshino N, et al. 2000. Exp. Anim. (Tokyo) 49:97. (FC)
- 12. Rout N, et al. 2010. PLoS One 5:e9787. (FC)

#### **Description:**

IL-2 is a potent lymphoid cell growth factor which exerts its biological activity primarily on T cells, promoting proliferation and maturation. Additionally, IL-2 has been found to stimulate growth and differentiation of B cells, NK cells, LAK cells, monocytes, and oligodendrocytes.

# Antigen References:

- 1. Fitzgerald K, *et al.* Eds. 2001. The Cytokine FactsBook. Academic Press, San Diego.
- 2. Taniguchi T, et al. 1993. Cell 73:5.
- 3. Nistico G. 1993. Prog. Neurobiol. 40:463.
- 4. Waldmann T, et al.