

PE/Cy7 anti-human CD197 (CCR7)

Catalog # / Size: 2366125 / 25 tests
2366130 / 100 tests

Clone: G043H7

Isotype: Mouse IgG2a, κ


Immunogen: CCR7-transfected cells

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific




Human peripheral blood lymphocytes were stained with CD3 FITC and CD197 (clone G043H7) PE/Cy7 (top) or mouse IgG2a, κ PE/Cy7 isotype control (bottom).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



Application References: 1. Kaczmarek Michaels K, *et al.* 2015. *J Immunol.* 194:3267. [PubMed](#)

Description: CCR7, also known as CD197, is a chemokine receptor that binds CCL19 and CCL21. CCR7 and its ligands link innate and adaptive immunity by affecting interactions between T cells and dendritic cells and their downstream effect. Naïve T cells enter the lymph node through high endothelial venules, which express CCL21. Dendritic cells and macrophages enter the lymph node through afferent lymphatics. The encounter of T cells and dendritic cells in the T cell zone is CCR7-dependent. In addition, during immunological surveillance, B cells recirculate between B-cell-rich compartments (follicles or B cell zones) in secondary lymphoid organs, surveying for antigen. After antigen binding, B cells move to the boundary of B and T zones to interact with T-helper cells; this B cell migration is directed by CCR7 and its ligands. CCR7-positive cancer cell expression has been associated with lymph node metastasis.

Antigen References: 1. Yanagihara S, *et al.* 1998. *J. Immunol.* 161:3096.
2. Charo IF, *et al.* 2006. *N. Engl. J. Med.* 354:610.
3. Reif K, *et al.* 2002. *Nature* 416:94.

4. Nakata B, *et al.* 2008. *O*