

Brilliant Violet 421™ anti-human CD186 (CXCR6)

Catalog # / Size: 2380065 / 25 tests
2380070 / 100 tests

Clone: K041E5

Isotype: Mouse IgG2a, κ


Immunogen: cDNA encoding human CXCR6

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 421™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 421™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Concentration: Lot-specific




Human peripheral blood mononuclear cells were stimulated with IL-2 for 9 days, and then stained with CD3 APC and CD186 (clone K041E5) Brilliant Violet 421™ (top) or mouse IgG2a, κ Brilliant Violet 421™ isotype control (bottom).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



Brilliant Violet 421™ excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by

U.S. Patent(s), pending patent applications and foreign equivalents.

Description: CD186, also known as CXCR6, is a 39 kD, G-protein coupled chemokine receptor, with seven transmembrane-spanning regions. CXCR6 is expressed on activated and memory T cells. Its ligand is the chemokine CXCL16. CXCR6 is involved in the recruitment of cells to inflamed sites and is also a coreceptor for HIV.

Antigen
References:

1. Kim CH, *et al.* 2001. *J. Clin. Invest.* 107:595.
2. Nanki T, *et al.* 2005. *Arthritis Rheum.* 52:3004.
3. Van der Voort R, *et al.* 2005. *Arthritis Rheum.* 52:1381.
4. Aust G, *et al.*